CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING



ACTIVE CONSTITUENTS: 96 g/L AZOXYSTROBIN 194 g/L TRITICONAZOLE

Dual Systemic Action to Control Over 10 Turf Diseases

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA Approval No.: 80146/133639HG



ACN: 117 986 615 43 Gap Road Sunbury Vic 3429 Ph: 1300 11 8873 (Ph: 1300 11 TURF) www.turfculture.com.au

Impala Fungicide Dual Action Systemic Fungicide

Impala Fungicide is an effective combination of two fungicides to prevent and control diseases in lawns. Impala's two active constituents work outside and inside the plant, helping prevent early fungal infection and established disease control, controlling all life cycle stages of turf diseases.

HOW TO USE:

Measure lawn area before application.

Situation	Disease	Rate	Critical Comments
Turf / Lawns	Leaf and Crown Diseases as listed; Anthracnose (Colletotrichum graminicola), Brown Patch (Rhizoctonia solani), Grey Leaf Spot (Pyricularia grisea), Helminthosporium Disease (Bipolaris spp, Drechslera spp, Exserohilum spp), Fusarium (Fusarium nivale, Microdochium nivale), Dollar Spot (Sclerotinia homoeocarpe), Pythium Leaf Blight,	Garden Sprayer Mix 60 mL in 5L of water and evenly spray onto 100 square metres of lawn Watering Can Mix 6 mL in 9 litres of water and evenly apply to 10 square metres of lawn	 For Preventative Control: Apply every 21-28 days when conditions favour disease development (e.g. warm, rainy weather). For Curative Control: Apply when disease symptoms first appear and repeat every 14-21 days. When spraying with Garden Sprayer ensure thorough and even coverage of the lawn. Irrigate the area sprayed with approx. 6 mm of rainfall equivalent commencing within 1 hour of application. DO NOT allow the product to dry on the leaf and crown surfaces. When irrigating the product into the soil after application DO NOT irrigate to the point of runoff (i.e. do not flood or irrigate more irrigation (mm) per hour than what the treated area's infiltration rate is – the soil moisture absorption rate). In high pressure Pythium curative situations use a suitable curative fungicide first and then follow up with Impala Fungicide.
	Seedling Damping Off (<i>Pythium spp</i>) <u>Root Diseases</u> as listed;	Garden Sprayer	When spraying with Garden Sprayer ensure thorough and even coverage of the lawn.
	Brown Patch (Rhizoctonia solani), Fairy Ring (Basidiomycetes fungi), Pythium Root Rot	Mix 60 mL in 5L of water and evenly spray onto 100 square metres of lawn	Irrigate the area sprayed with approx. 6 mm of rainfall equivalent commencing within 1 hour of application. DO NOT allow the product to dry on the leaf and crown surfaces. DO NOT irrigate to the point of runoff (i.e. DO NOT flood or irrigate more irrigation (mm) per hour than what the treated area's infiltration rate is – the soil moisture absorption rate).
	(<i>Pythium spp</i>), Ectotrophic Root Infecting Fungi (ERI) including;	Watering Can Mix 6 mL in 9 litres of water and evenly	Irrigate the area sprayed again with approx. 6 mm of rainfall equivalent 3 days after application. Fairy Ring: Apply preventatively when conditions are
	Couchgrass Decline (Gaeumannomyces graminis var. graminis), Spring Dead Spot (Ophiosphaerella narmari),	apply to 10 square metres of lawn	favorable for disease development or as an early curative as soon as possible after first symptoms are detected. Make a second application 21 days later. Ectotrophic Root Infecting Fungi (ERI) - Autumn strategy Preventative applications: Spray in late February. Make a second application 1 month later.
	Take-all Patch (Gaeumannomyces graminis var. avenae)		Ectotrophic Root Infecting Fungi (ERI) - Spring and summer strategy Preventative applications: Spray in mid-October. Make a second application 1 month later. Early curative applications: Spray as soon as possible after first symptoms are detected. Make a second application 1 month later.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

HOW TO PREPARE

Prior to pouring, shake container vigorously, then add the required quantity of Impala Fungicide to clean water in a garden sprayer (or watering can) and mix thoroughly. The prepared spray must be used on the day it is mixed.

DO NOT apply to waterlogged soil. DO NOT apply if heavy rains or storms that are likely to cause runoff are forecast within 3 days. DO NOT apply when soil is dry, and plants are suffering from moisture stress. DO NOT spray when shade temperatures are likely to exceed 30°C. DO NOT use on edible plants.

Compatibility

DO NOT mix with products or fertiliser with high salt content, i.e. ferrous sulphate or ammonium sulphate.

PRECAUTIONS

DO NOT allow entry into treated areas until the spray has dried (including pets). DO NOT spray Impala Fungicide where spray drift may reach apple trees. Impala Fungicide is extremely phytotoxic to certain apple varieties.

Apply no more than 2 consecutive applications of Impala Fungicide per disease per season. DO NOT allow containers or spray to get into drains, sewers, ponds or streams. DO NOT feed grass clippings from treated areas to poultry or other animals.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated, dry place out of reach of children and pets. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID INSTRUCTIONS

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

CONDITIONS OF SALE

The use of Impala Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by Turf Culture regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Turf Culture accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

In a Transport Emergency Dial 000 Police or Fire Brigade